

Abstract

The F_2 structure functions of the inelastic lepton-hadron scattering is calculated in the case of non-zero intermediate gluon-quarks self-energy M_{gq}^2 and quasielastic limit. It is shown that in the quasielastic limit the evolution equation can be solved in case massive particles as QCD ladder calculation results. The intermediate gluon-quarks self-energy, or polarization mass is introduced as a parameter in two cases: 1) not momentum dependence and 2) momentum dependence cases. At the case 1) of massive particles we compare our result with Shirkov calculates (D.V.Shirkov 1999). It has been shown (case 2.) that the proton structure function F_2 calculated at realistic values of M_{gq}^2 in high energy limit transfer to standard QCD ladder calculation results.

PACS : 13.85.Fb, 14.70.D

Keywords : Structure functions, Deep Inelastic Scattering, Quasielastic Scattering, Total Cross Section, Gluon Self-Energy, Gluon mass.

On the theory of quasielastic lepton-hadron scattering

Ya.D. Krivenko-Emetov *

*Institute for Nuclear Research, Ukrainian Academy of Sciences,
Prospect Nauki, 47, Kiev 03680, Ukraine
Applied Physics Department of Kiev National Technical University,
Prospect Pobedu, 37, Kiev 03056, Ukraine*

February 1, 2008

1 INTRODUCTION

It is known that gluons are massless particles due to 1) principles renormalization theory, 2) asymptotic freedom and 3) exact $SU(3)_c$ symmetry and local invariance assumption. However, the question of a non-zero gluon mass possibility and its physical consequences is discussed for a long time in literature. It was assumed, in particular, that gluons become massive due to symmetry breaking [1, 2, 3]. In this case the requirements 1) and 2) can be fulfilled. In Section 2, proposed another mechanism for the gluon mass generation (of course our supposition does not extenuate the significance of result [1]). Gluon becomes massive due to the interaction with quarks. In this case the quarks-gluons mass is nothing else but polarization correction to its self-energy. This quantity is a real correction to quarks-gluon self energy, no matter if particle is on or out of mass surface. Therefore, this assumption does not lead to impossibility to use Katkowski rules for the full cross section calculations by using the optical theorem [4]. We assume that this quantity at low energy 1.) has not energy momentum dependence and is an effective theory parameter M^2 , which can have an arbitrary sign, 2.) leads as $k^2 L^2(k^2)$, where $L^2(k^2)$ is k^2 momentum dependence and k^2 -transfer moment of gluon and 2.1) use parametrization [5] put $L^2 = \tilde{L}^2(\ln[(\Lambda_N^2 + k^2)/\mu_{QCD}^2]/\ln[\Lambda_N^2/\mu_{QCD}^2])$ ($\ln(\mu^2) = \ln(\Lambda_{QCD}^2) + 4\pi/(\beta_0\alpha_0)$), where Λ_N - uncalculable factor that describe unperturbative contribution in strong coupling or quarks-gluon self-energy, and $\Lambda_{QCD} \simeq 216(\pm 25) MeV$ [9] is the constant of QCD. In our model M_{gq}^2 not

*E-mail address: yakr@kinr.kiev.ua

only gluon self-energy. This parameter also discount (check) quarks self-energy according theory of renormalizable (see eq. (1) andless). Its calculation can not be performed exactly since the perturbation theory is not applicable in the QCD theory. Justification for investigate this not momentum dependence parameter can be obtained in modern Effective Field Theory and Quantum Hadrodynamics (see for example [6],[7],[8]). This correction M_{gq} must be small, but greater with respect to quark mass. However, it can lead to some physical effects. Namely, we show that the self-energy correction leads to a significant change in the lepton-hadron scattering cross section. This effect takes place because the set of ladder diagrams with infinite number of gluon lines must be taken into account. Thus, the intercept correction is of order M_{gq}^2/Q^2 , and the logarithm is of order $M_{gq}^2(Q^2, \Lambda_{QCD}^2)$, where Q^2 is the squire of transferred momentum ($Q^2 = -q^2$). Formally, the contribution of structural function to cross section becomes important at small Q^2 , but at these moment perturbative ladder approximation does not work, and the polarization correction is not known so far. It is known that at large transfer moment polarizability contribution can be taken into account by renormalization of the coupling constant $g \rightarrow g(k^2)$, since these contributions can be factorized and by using the Ward identity [10]. In this case "polarization mass" is

$$M_{pol}^2(k^2) \simeq_{k^2 > Q_0^2} k^2 \tilde{L}^2 (\ln[(\Lambda_N^2 + k^2)/\mu_{QCD}^2] / \ln[\Lambda_N^2/\mu_{QCD}^2])^2$$

$$\rightarrow_{k^2 \rightarrow \infty} -k^2 b \ln(k^2/\mu^2), \quad (\tilde{L}^2 / \ln(\Lambda_N^2/\mu_{QCD}^2) \sim -b) \quad (1)$$

$(Im[\frac{e}{1+e \ln(k^2/\mu^2)} \frac{1}{k^2+i0}] = e\delta(k^2 + k^2 e \ln(k^2/\mu^2)) = e\delta(k^2 - M^2) = \frac{e}{1+e \ln(k^2/\mu^2)} \delta(k^2))$ where $b = 11 - (2/3)n_f, n_f$ -quarks flowers [21]. Comparison of the cross-section calculated in this way with experimental results is a possible indirect way to estimate the value of possible quarks-gluon mass [1, 2, 11, 12, 13]. It is important to say that processes of creation of other particles also give a contribution to "gluon mass", therefore our effective M_g is assumed to take into account these contributions too. This gives as a possibility to describe the quasiperturbative region of scattering when Q^2 is not very large, and maybe even when Q^2 is rather large, in contrast to the previous results, we the polarization correction was calculated at large Q^2 [14, 15, 16, 17, 18, 19]. It is shown (eq.(16)), that in case large Q^2 our results traversed to results [14, 15, 16, 17, 18, 19]. At fig1.,2 we present the model [1] (14), and our estimation with massive effective particles with different value of parameter and variables. Of course, this solution (14),(15) is not correct at very small Q^2 , when one loop approximation is incorrect. But the one-loop solutions at large Q^2 and our solution (14), (16) at intermediate Q^2 will be "sewed" at some point of transfer momentum.(method "sewed" see for example [20],[5]). Analytically this "sewed" give formula (16)-(18). Also, the non-singlet diagrams give the biggest contribution at $x \rightarrow 1$, but in this case delta-functions will give large contributions in terms which is difficult to calculate, and we did not take into account. It was shown in [22],

for example, that these diagrams lead to a good description of experiment at $x > 0.65$. Finally, Section 3. contains a summary and conclusions.

2 ONE-LOOP DIAGRAMS

Let us use the axial gauge $A_\mu \eta^\mu = 0$. It should be mentioned, that in the case of massive particles the gauge is not arbitrary and the propagator is

$$G_{\mu\nu} = -\frac{1}{p^2 - m^2 + i0} \left(g_{\mu\nu} - \frac{p_\mu p_\nu}{m^2} \right).$$

As it was mentioned in the Introduction, we assume that virtual gluons have non-zero self-energy, or polarization mass $M_g(Q^2)$. In this case the hadron tensor has the next form [23] ($S_2(V)$ -color multiply, M-mass of hadron)

$$W_{\mu\nu}(p, q) = \frac{g^2 S_2(V)}{2M} \int \frac{d^4 k}{(2\pi)^3} \times \frac{\delta[(p-k)^2 + M_{gq}^2] \delta[(q+k)^2]}{k^4} d^{\rho\sigma} (p-k) T_{\rho\mu\nu\sigma} + \Delta W_{\mu\nu}. \quad (2)$$

Where $\Delta W_{\mu\nu}$ discount, that M_{gq}^2 can have more composite structure, then constant or $k^2 L^2$. Here we substitute the quark and gluon propagator by delta-functions with arguments equal to the denominator of propagator expressions.

Let us introduce the Sudakov variables:

$$k_\mu = \xi p_\mu + \beta q'_\mu + k_{\perp\mu}$$

with

$$q'^2 \simeq p^2 \simeq 0, \quad pq' \simeq pq,$$

and a new variable x : $q' = q + xp$. In this case the integral in (2) can be transformed as

$$\begin{aligned} & \int d^4 k \delta[(p-k)^2 - M_g^2] \delta[(q+k)^2] = \\ & = 2\pi pq \int d\beta d\xi dk_\perp^2 \frac{1}{2pq(1-\xi)} \delta\left[\beta - \frac{k_\perp^2 - M_{gq}^2}{2\pi pq(1-\xi)}\right] \\ & \times \frac{1}{2pq} \delta\left[\xi - x + \frac{(1-x)k^2}{2pq} + \frac{M^2 x}{2pq}\right] \end{aligned} \quad (3)$$

(see Appendix) where we use the substitution $d^4 k = 2\pi pq d\beta d\xi dk_\perp^2$ and approximation $-\frac{k}{q'} + \beta^2 \frac{q'^2}{2pq(1-\xi)} \rightarrow 0$.

If

$$\beta < 1 - \xi \Rightarrow \xi \sim x \quad (4)$$

,where

$$\xi \simeq x_f \simeq \frac{2x_b}{1 + \sqrt{1 + 4M_b^2 x_b^2 / Q^2}} \quad (5)$$

([24],[22], when $Q^2 \gg M_n^2$, $\xi \rightarrow x_b = x$). Similarly to [14, 17], the appearance of the term $\ln(-q^2/p^2)$ is related to the phase space regions, where the relations (4) are correct. Therefore, variable x which corresponds to the parton model variable ia part of longitudinal momentum and it justifies the approximation $q'^2 \simeq 0$.

The square of k_μ can be expressed as $k^2 = \xi^2 p^2 + \beta \xi p^\mu q'_\mu + \xi p^\mu k_{\perp,\mu} + \beta \xi q'^\mu p_\mu + \beta^2 q'^2 + \beta q'^\mu k_{\perp,\mu} + \xi k_{\perp}^\mu p_\mu + \beta k_{\perp,\mu} q'^\mu + k_\perp^2 \simeq 2\beta \xi(q'p) + k_\perp^2$ use (3), obtain

$$k^2 \simeq \frac{k_\perp^2}{(1-\xi)} - \frac{M_{gq}^2 \xi}{(1-\xi)}. \quad (6)$$

Convolution of the hadron and gluon polarization tensors with putting $\eta = q'$ and using of γ -matrix algebra, (3),(6) gives the main contribution to the hadron tensor, which corresponds to the tree diagram. If $\beta < 1 - \xi \Rightarrow \xi \sim x$, obtain

$$\begin{aligned} d^{\rho\sigma}(p-k)T_{\rho\mu\nu\sigma} - \Delta W_{\mu\nu} &= 2tr[\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu](-\hat{k}\hat{p}\hat{k} + \\ &+ \frac{k^2}{(p-k, q')}[(pq')\hat{k} + (kq')\hat{p} + (p-k, p)\hat{q}']) \end{aligned} \quad (7)$$

and use (20)-(23)(see Appendix), obtain at case

1.1) $M^2(k^2) \simeq M^2$

$$\simeq 2k^2 tr[\hat{p}\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \left(\frac{1+\xi^2}{1-\xi} + \frac{M_g^2 \xi}{pq} + \frac{M_g^2 \xi}{pq(1-\xi)} - \frac{M_g^4 \xi}{2pqk^2} \right) \quad (8)$$

2.1) and at case $M^2(k^2) \simeq L^2 k^2$ (supply it at (23),obtain)

$$\simeq 2k^2 tr[\hat{p}\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \left(\frac{1+\xi^2}{1-\xi} + L^2 \xi \right) \quad (9)$$

and if $\beta > 1 - \xi \Rightarrow \xi \sim x - \frac{(1-x)k^2}{2pq} - \frac{M^2 x}{2pq}$ (see Appendix, eq.(24))

$$\begin{aligned} d^{\rho\sigma}(p-k)T_{\rho\mu\nu\sigma} - \Delta W_{\mu\nu} &\simeq 2tr[\hat{p}\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \\ &\left(-\frac{(k^2 - M_g^2)^2 \xi}{2pq} + M_g^2 \xi + k^2(1-x) + \frac{k^2 2\xi}{\beta} - \frac{k^2 2\xi(1-\xi)}{\beta} \right) \end{aligned} \quad (10)$$

2.2.) if $M^2 = k^2 L^2$, $\beta > 1 - \xi \Rightarrow \xi \sim x - \frac{(1-x)k^2}{2pq} - \frac{k^2 L^2 x}{2pq}$ (see Appendix, eq. (25))

$$(eq.10) \simeq 2k^2 tr[\hat{p}\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \left(L^2 x + (1-x) - \frac{2(x - x^2 + L^2 x^2)}{(1-L^2)} \right) \quad (11)$$

To obtain the expression (8),(10),(11) the first and the second terms (equal to $(1-\xi)k^2$ and $2\xi k^2/(1-\xi)$, correspondingly) in gluon propagator were considered. We also neglecting $1/k^4$ and k^4 , because at high k^2 $|M_g^2| \simeq k^2 \ln(k^2/\Lambda_{QCD}^2) > k^2$, and $k^2/2pq = k^2 x/Q^2 \simeq k^2 L^2/Q_{Q^2 \rightarrow \infty}^2 \rightarrow 0$. However, in contrast to the paper [14, 17], first and the third term in (22) was also taken into account, since it gives the contribution from the polarization mass in (8) and (9) respectively. If $\beta < 1 - \xi \Rightarrow \xi \sim x$ use (3),(6),(8), obtain in cases
1.1.) $M^2(k^2) \simeq M^2$

$$(26) = \frac{3g^2 S_2(V)}{2M_n(2\pi)^3} \pi \int_{m^2 - f(M^2)}^{Q^2 - f(M^2)} \frac{d\beta d\xi dk^2}{k^2} 2tr[\hat{p}\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \times \\ \times \left(\frac{1+\xi^2}{1-\xi} + \frac{M_g^2 \xi}{pq} + \frac{M_g^2 \xi}{pq(1-\xi)} - \frac{M_g^4 \xi}{2pqk^2} \right) \times \\ \times \frac{1}{2pq} \delta[\beta - (k^2 - M_g^2)/2pq] \delta((\xi - x) + \frac{(1-x)k^2}{2pq} + \frac{M^2(x - \xi^2)}{2pq}) \quad (12)$$

2.1.) According (1), put $M^2 \simeq k^2 L^2$

$$= \frac{3g^2 S_2(V)}{2M_n(2\pi)^3 2pq} \pi \ln \left[\frac{Q^2 - Q^2 L^2 x/(1-x)}{m^2 - m^2 L^2 x/(1-x)} \right] 2[8\xi p_\nu p_\mu - \\ 4M\nu g_{\mu\nu} + \dots] \left(\frac{1+x^2}{1-x} + A^2 x + \frac{A^2 x}{1-x} \left(\frac{1+x^2}{1-x} + A^2 x \right) \right) \quad (13)$$

According (12) we obtain with equations (2) and (8) give (if $\beta < 1 - \xi$, $M^2(k^2) = M^2$)

$$f(x, Q^2) - \Delta f(x, Q^2) \sim g^2 \frac{M_g^4 x^2}{Q^4} + \\ + g^2 \ln \left[\frac{Q^2 - M_g^2 x/(1-x)}{m^2 - M_g^2 x/(1-x)} \right] \left(\frac{1+x^2}{1-x} + \frac{2M_g^2 x^2}{Q^2} + \frac{2M_g^2 x^2}{Q^2(1-x)} \right) \quad (14)$$

(for massive gluons $\Delta f(x, Q^2) = 0$ and we have put $x = Q^2/2pq$ in the last expression), and if $M^2 = k^2 L^2$, $\beta < 1 - \xi$ (13) give (put $L^2 = \tilde{L}^2 (\ln[(\Lambda_N^2 + Q^2)/\Lambda_{QCD}^2]/\ln[\Lambda_N^2/\Lambda_{QCD}^2])$, $|\ln(\Lambda_{QCD}^2)| >> 4\pi/(\beta_0 \alpha_0)$)

$$f(x, Q^2) - \Delta f(x, Q^2) \sim g^2 \ln \left[\frac{Q^2 - Q^2 L^2 x/(1-x)}{m^2 - m^2 L^2 x/(1-x)} \right] \\ \left(\frac{1+x^2}{1-x} + A^2 x + \frac{A^4 x^2}{(1-x)} + \frac{A^2 x(1+x^2)}{(1-x)^2} \right) \rightarrow_{Q^2 \rightarrow \infty} \quad (15)$$

$$\begin{aligned} & \rightarrow_{Q^2 \rightarrow \infty} g^2 \left(\ln(Q^2/m^2) + \ln \left[\frac{Ln(Q^2/\Lambda_{QCD}^2)}{Ln(m^2/\Lambda_{QCD}^2)} \right] \right) \times \left(\frac{1+x^2}{1-x} + \right. \\ & \left. \tilde{A}^2 \left(\frac{x(1+x^2)}{(1-x)^2} + x + \frac{\tilde{A}^2 x^2}{(1-x)} \right) \right) \end{aligned} \quad (16)$$

Compare this result with standard result at high Q^2 , obtain that at $Q^2 \rightarrow \infty$

$$\begin{aligned} \Delta f(x, Q^2) & \simeq -g^2 \ln(Q^2/m^2) \left[\left(\frac{1+x^2}{1-x} \right) + 1 + \frac{\tilde{A}^2}{g^2 \ln(Q^2/m^2)} \right. \\ & \left. \times \ln \left[\frac{Ln(Q^2/\Lambda_{QCD}^2)}{Ln(m^2/\Lambda_{QCD}^2)} \right] \right] \left(\frac{x(1+x^2)}{(1-x)^2} + x + \frac{\tilde{A}^2 x^2}{(1-x)} \right) \end{aligned} \quad (17)$$

and at $L^2 \rightarrow 0$ $\Delta f(x, Q^2) \simeq 0$ But this result (eq. (16)) give badly behavior at $L \rightarrow 0$ ($A^2 \rightarrow 0$), because at $Q^2 + \Lambda_N^2 = \Lambda_{QCD}^2$ must be fulfillment condition $m^2 = \Lambda_{QCD}^2 - \Lambda_N^2$. Good behavior have another renormalizable result, that received similarly (we know polarized mass only at high k^2 . This unknown function we can transform into Teilor series $M^2 = const_+^-(Q^2/Q_{unpert}^2) f(m^2)_+^- \dots \rightarrow_{Q^2 > Q_{unpert}^2} Q^2 \ln(m^2/\Lambda_{QCD}^2), Q_{unpert}^2 \sim 3 - 5 Gev^2$, L^2 gives (1)):

$$\begin{aligned} f(x, Q^2) & \simeq const * \frac{48g^2 S_2(V)}{2(2\pi)^3} \pi \left[\ln \left[\frac{Q^2 - Q^2 L^2(Q^2) x / (1-x)}{m^2 - Q^2 f(\tilde{m}^2) x / (1-x)} \right] \right] \times \\ & \times \left(\frac{1+x^2}{1-x} + A^2 x + \frac{A^2 x}{1-x} \left(\frac{1+x^2}{1-x} + A^2 x \right) \right) - \ln \left[\frac{Ln(Q^2/\Lambda_{QCD}^2)}{Ln(m^2/\Lambda_{QCD}^2)} \right] \times \\ & \times \tilde{A}^2 \left(x + \frac{x}{1-x} \left(\frac{1+x^2}{1-x} + \tilde{A}^2 x \right) \right) \end{aligned} \quad (18)$$

3 CONCLUSIONS

It has been shown that the contribution to inelastic lepton-hadron scattering cross section from the quarks-gluon polarization mass can be rather large. Polarization mass consideration can be a possible way to decrease the difference between experimental data and theoretical calculations of the structure functions of inelastic hadron lepton scattering. Another application of the results obtained in this paper is a possible estimation of the quarks-gluon self-energy results calculated by formula (14) (18), which contains quarks-gluon self-energy as a function-parameter. We believe that quarks-gluon polarization mass presence will make the condition $k_1 \ll k_2 \ll \dots \ll k_N$ weaker [25], even if these other processes are not taken into account. And that this gives as a possibility to describe the quasiperturbative region of scattering when Q^2 is not very large, and

maybe even when Q^2 is rather large, in contrast to the previous results, where the polarization correction was calculated at large Q^2 [14, 15, 16, 17, 18, 19]. At fig.1.,2. we present our calculation for the model [1] (14), and our estimation (18) with effective polarization mass with different value of parameter and variables. How apparently at fig.1 "skailing" increase with increase mass of gluons. In fig.2 "skailing" is more visible and in principle can be used for describe experimental data.

Acknowledgments

Ya.D.K.-E. would like to thank V.M. Turkowski, A.P.Kobushkin and V.V.Davidovsky for useful discussions which have resulted in an improvement of the paper. Ya.D.K.-E. acknowledges the financial support from the Applied Physics Department of Kiev National Technical University, Project N2700/162.He also thanks V.M.Kolomic for aid.

APPENDIX

For obtain (3), use

$$\begin{aligned} \delta((p-k)^2 - M_g^2) &= \delta(\beta^2 q'^2 + k_\perp^2 - 2pq'\beta(1-\xi) - 2pk_\perp(1-\xi) - M_{gk}^2) \simeq \frac{1}{2pq(1-\xi)} \delta(-\beta + \\ &\frac{k_\perp^2 - M_g^2}{2\pi pq(1-\xi)} - \frac{k_\perp}{q'} + \beta^2 \frac{q'^2}{2pq(1-\xi)}) \simeq \frac{1}{2pq(1-\xi)} \delta(-\beta + \frac{k_\perp^2 - M_g^2}{2\pi pq(1-\xi)}), \text{ and} \\ \delta(k+q)^2 &\simeq \delta((\beta+1)q' + (\xi-x)p + k_\perp)^2 \simeq \delta(2(\beta+1)(\xi-x)pq + k_\perp^2) \simeq \\ &\frac{1}{2pq} \delta((\xi-x) - \beta x + \frac{k^2}{2pq} - \frac{k^2\xi}{2pq} + \beta\xi + \frac{M_g^2\xi(1-\xi)}{2pq}) \simeq \frac{1}{2pq} \delta((\xi-x) + \frac{(1-x)k^2}{2pq} + \frac{M^2(x-\xi^2)}{2pq}) \\ &\simeq \frac{1}{2pq} \delta(\xi-x + \frac{(1-x)k^2}{2pq} + \frac{M^2x}{2pq}) = \frac{1}{2pq} \delta(\xi-x + \frac{(1-x)k^2x}{Q^2} + \frac{M^2x^2}{Q^2}) \end{aligned} \quad (19)$$

Supplied

$$-(p - (p - k))^2 = -p^2 - (p - k)^2 + 2p(p - k) = -p^2 - M_g^2 + 2p(p - k) \quad (20)$$

and $(p - k)^2 = M_g^2 \rightarrow p(p - k) = p^2/2 - k^2/2 + M_g^2/2 \simeq p^2/2 + M_g^2/(2(1 - \xi)) - k_\perp^2/(1 - \xi)$ in to (7) , obtain

$$\begin{aligned} d^{\rho\sigma}(p - k)T_{\rho\mu\nu\sigma} &= 2tr[\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \\ &\times (-\hat{k}\hat{p}\hat{k} + \frac{k^2}{(p - k, q')}[(pq')\hat{k} + (kq')\hat{p} + (p^2/2 + M_g^2/2 - k^2/2)\hat{q}']) \end{aligned} \quad (21)$$

and use

$$\hat{k}\hat{p}\hat{k} = 2\beta(pq)(\beta q' + k_\perp) - (k^2(1 - \xi) + M_g^2\xi)\hat{p} \quad (22)$$

$(p-k)q' = (1-\xi+\beta)\frac{k^2-k_\perp^2}{2\beta\xi}, \quad (pq')\hat{k} + (kq')\hat{p} = \frac{k^2-k_\perp^2}{2\beta\xi}(\hat{k} + \xi\hat{p}),$
 $k^2 = 2\beta\xi(pq) + k_\perp^2, \quad \beta \simeq (k_\perp^2 - M_g^2)/(2pq(1-\xi)) = (k^2 - M_g^2)/2pq$ rewrite (21)
 as

$$\begin{aligned}
 &\simeq 2tr[...](-2\beta(pq)\beta\hat{q}' + (k^2(1-\xi) + M_g^2\xi)\hat{p} + \frac{k^2}{1-\xi+\beta}(\hat{k} + \xi\hat{p} - \frac{k^2 - M_g^2}{2pq}\hat{q}')) \\
 &\simeq 2tr[...]\left(-2\beta(pq)\beta\hat{q}' + (k^2(1-\xi) + M_g^2\xi)\hat{p} + \frac{k^2}{1-\xi}(\beta\hat{q}' + 2\xi\hat{p} - \beta\hat{q}') - \frac{k^2 2\beta\xi\hat{p}}{1-\xi} \right) \\
 &\simeq 2tr[...]\left(k^2\frac{1+\xi^2}{1-\xi} - \frac{(k^2 - M_g^2)^2\xi}{2pq} - \frac{(k^2 - M_g^2)k^2\xi}{pq(1-\xi)} + M_g^2\xi \right) \simeq (eq. 8)(23)
 \end{aligned}$$

if $\beta < 1 - \xi$, and if $\beta > 1 - \xi$, obtain

$$\begin{aligned}
 &\simeq 2tr[...](-2\beta(pq)\beta\hat{q}' + (k^2(1-\xi) + M_g^2\xi)\hat{p} + \frac{k^2}{1-\xi+\beta}2\xi\hat{p}) \\
 &\simeq 2tr[\hat{p}\gamma_\mu(\hat{q}+\hat{k})\gamma_\nu]\left(-\frac{(k^2 - M_g^2)^2\xi}{2pq} + M_g^2\xi + k^2(1-\xi) + \frac{k^2 2\xi}{1-\xi+\beta} \right) = (eq.(10))(24)
 \end{aligned}$$

1.) if $M^2 = k^2 L^2$, $\beta > 1 - \xi$

$$\simeq 2tr[\hat{p}\gamma_\mu(\hat{q}+\hat{k})\gamma_\nu]\left(k^2 L^2 x + k^2(1-x) + \frac{2\xi Q^2}{(1-L^2)x} - \frac{2\xi(1-\xi)Q^2}{(1-L^2)x} \right) = (eq.(11))(25)$$

At the case $M^2(k^2) \sim M^2$, and $\beta < 1 - \xi$ supply (3),(8) and (6) into (2), obtain

$$\begin{aligned}
 &\frac{3g^3}{2M_n(2\pi)^3} 2\pi pq \int \frac{d\beta d\xi dk_\perp^2}{k^4} 2k^2 tr[\hat{p}\gamma_\mu(\hat{q}+\hat{k})\gamma_\nu] \left(\frac{1+\xi^2}{1-\xi} + \frac{M_g^2\xi}{pq} + \frac{M_g^2\xi}{pq(1-\xi)} - \frac{M_g^4\xi}{2pqk^2} \right) \\
 &\quad \frac{1}{2pq(1-\xi)}\delta[\beta - \frac{k_\perp^2 - M_g^2}{2\pi pq(1-\xi)}] \times \frac{1}{2pq}\delta((\xi-x) + \frac{(1-x)k^2}{2pq} + \frac{M^2(x-\xi^2)}{2pq}) \\
 &= \frac{3g^3}{2M_n(2\pi)^3} \pi \int_{m^2-f(M^2)}^{Q^2-f(M^2)} \frac{d\beta d\xi d(k^2 + \frac{M^2\xi}{1-\xi})}{k^2} 2tr[\hat{p}\gamma_\mu(\hat{q}+\hat{k})\gamma_\nu] \left(\frac{1+\xi^2}{1-\xi} + \frac{M_g^2\xi}{pq} + \frac{M_g^2\xi}{pq(1-\xi)} - \frac{M_g^4\xi}{2pqk^2} \right) \\
 &\quad \frac{1}{2pq}\delta[\beta - (k^2 - M_g^2)/2pq]\delta((\xi-x) + \frac{(1-x)k^2}{2pq} + \frac{M^2(x-\xi^2)}{2pq}) \quad (26)
 \end{aligned}$$

2.) According (1), put $M^2 \simeq k^2 L^2$ into (2) and use (9), obtain

$$= \frac{3g^3}{2M_n(2\pi)^3} \pi \int_{m^2-f(M^2, Q^2)}^{Q^2-f(M^2, Q^2)} \frac{d\beta d\xi d(k^2 + k^2 \frac{L^2\xi}{1-\xi})}{k^2} 2tr[\hat{p}\gamma_\mu(\hat{q}+\hat{k})\gamma_\nu] \left(\frac{1+\xi^2}{1-\xi} + L^2\xi \right)$$

$$\begin{aligned}
& \frac{1}{2pq} \delta[\beta - (k^2 - M_g^2)/2pq] \delta((\xi - x) + \frac{(1-x)k^2}{2pq} + \frac{M^2(x - \xi^2)}{2pq}) \\
&= \frac{3g^3}{2M_n(2\pi)^3} \pi \int_{m^2 - m^2 L^2(m^2)\xi/(1-\xi)}^{Q^2 - Q^2 L^2(Q^2)\xi/(1-\xi)} \frac{d\beta d\xi dk^2 (1 + \frac{L^2 \xi}{1-\xi})}{k^2} 2tr[\hat{p}\gamma_\mu(\hat{q} + \hat{k})\gamma_\nu] \left(\frac{1 + \xi^2}{1 - \xi} + L^2 \xi \right) \\
& \frac{1}{2pq} \delta[\beta - (k^2 - M_g^2)/2pq] \delta((\xi - x) + \frac{(1-x)k^2}{2pq} + \frac{M^2(x - \xi^2)}{2pq}) \\
& tr[\hat{p}\gamma_\mu(\hat{q} + \xi\hat{p})\gamma_\nu] = 4[p_\mu(\xi p + q)_\nu + (\xi p + q)_\mu p_\nu - p(\xi p + q)g_{\mu\nu}] = [8\xi p_\nu p_\mu - \\
& 4M\nu g_{\mu\nu} + \dots] \\
& \sim \frac{24g^2 S_2(V)\pi}{M_n(2\pi)^3} \ln \left[\frac{Q^2 - Q^2 L^2 x/(1-x)}{m^2 - m^2 L^2 x/(1-x)} \right] \left(1 + \frac{A^2 x}{(1-x)} \right) \\
& [\xi \frac{p_\nu p_\mu}{8 * 2pq} - \frac{2M\nu g_{\mu\nu}}{4 * 2pq} + \dots] \left(\frac{1 + x^2}{1 - x} + A^2 x \right) = (13) \tag{27}
\end{aligned}$$

where $A^2 \sim -b \ln \left[\frac{\Lambda_n^2 + Q^2 - Q^2 L^2 x/(1-x)}{\Lambda_n^2 + m^2 - m^2 L^2 x/(1-x)} \right] \xrightarrow{Q^2 \rightarrow \infty} -b \ln \left[\frac{Ln(Q^2/\Lambda_{QCD}^2)}{Ln(m^2/\Lambda_{QCD}^2)} \right] = \tilde{A}^2$

References

- [1] D.V. Shirkov, *Phys. of Atom. Nucl.* **62** (1999) 2082.
- [2] K. Cahill, **hep-ph/9812312**.
- [3] N.F.Nelipa, *Physics of elementary particle. Gauge fields.*(Russ.),Nuke, Moscow, 1985
- [4] E.M. Lifshitz et al, *Quantum Electrodynamics*, Nuke, Moscow, 1989.
- [5] See, for example M.F.Gari and W.Krumpelmann, *Phys.Lett***B274**,159(1992); E.Lomon *nucl-th/0203081* (2002); Ya.D.Krivenko-Emetov, A.P.Kobushkin *in press*(2003)
- [6] J.D. Walecka, *Theoretical Nuclear and Subnuclear Physics* (Oxford Univ. Press, 1995)
- [7] S. Weinberg, *Phys. Lett.* **B251**, 288 (1990)
- [8] S. Weinberg, *Nucl. Phys.* **B363**, 3 (1992)
- [9] K.Hagiwara et al., *Phys. Rev* **D66**, 010001 (2002)
- [10] See, for example N.N.Bogolubov,D.V.Shirkov, *Introduction in Theory Gauge Fields* (Russ.), Moscow ,1976

- [11] D.V. Shirkov, *Nucl. Phys.* **B 371** (1992) 467.
- [12] Yu.L. Dokshitzer, D.V. Shirkov, *Z. Phys.* **C 67** (1995) 449.
- [13] D.V. Shirkov, S.V. Mikhailov, *Z. Phys.* **C63** (1995) 467.
- [14] V.N. Gribov, L.N. Lipatov, *Yad. Fiz.* **15** (1972) 781 and 1218
[*Sov. J. Nucl. Phys.* **15** (1972) 438 and 675] and G.Altarelli G.Parisi *Nucl. Phys.*, **B 126** 298 (1977)
- [15] Y.Y.Balitskiy and L.N.Lipatov, *Sov. Phys. JETP* **28**, 882 (1978);
E.A.Kuraev, L.N.Lipatov, and V.S.Fadin, **45**,199 (1977)
- [16] D.J. Gross, *Phys. Rev. Lett.* **32** (1974) 1071.
- [17] Yu.L. Dokshitzer, D.I. Dyakonov, S.I. Troyan, *Phys. Rep.* **58** (1980) 270.
- [18] E. Kuraev, B. Fadin , *Yad. Fiz.* **41**, (1985) 733 [*Sov. J. Nucl.Phys.* **41**, (1985) 733].
- [19] I. Akushevich,E. Kuraev,B. Shaikhatdenov *Phys. of Element. Part. and Atomic Nuclei.*, (Russ.), **32** N3 491 (2001).
- [20] A.P.Kobushkin and Ya.D.Krivenko-Emetov **nucl-th/0112009**. (2001)
(eq. (22) of [20] must be multiply at a faktor $\alpha_s^4(Q^2)$.)
- [21] F.J.Yndurain *Quantum Chromodynamics*,
Springer-Verlag,N.-Y.,Berlin,Tokyo (1983)
- [22] S.R.Kelner,D.A.Timashkov, *Phys. of Atomic Nuclei* **64** N9 (2001) 1802.
- [23] See, for example Cheng, Li, *Gauge Theory of Elementary Particle Physics*,
Clarendon Press, Oxford University Press (1994).
- [24] H.Georgi and H.D.Politzer, *Phys. Rev.* **D.14**, 1829 (1976); *Phys.Rev.Lett.* **36**, 1281 (1976); A. de Rujula et al.,*Ann. Phys.(N.Y.)* 103, 315 (1977)
- [25] Ya.D.Krivenko-Emetov *in press*.

Figure caption

Fig.1 Our calculation for structural function F_2 in Shirkov approximation [1] (14) is presented as a function of Q^2 at different values of sign of the effective massive particles square M_p^2 and in the case of massless particles (16)-(17).

Fig.2 Structural function F_2 in our approximation (18) is presented as a function of Q^2 at different values of parameters of the effective "polarized masse" $M_g^2(\Lambda_N^2 = \Lambda = 3\text{GeV}^2)$. This results compare with well known perturbative result (16)-(17) for massless gluons.



